



If gardens are a window on our mind,

I see a passion for the European environment that indicates that we are still, at heart, uncomfortable in our own land.

If we see no beauty in Australian natives... can we really count ourselves as having a truly sustainable future adapted to

Australian conditions? Tim Flannery


## Increasing temperatures



TIME SERIES: 1884 TO 2015

Data source: NASA/GISS

Credit: NASA Scientific Visualization Studio




TIME SERIES: 1884 TO 2015

Data source: NASA/GISS

1990


Credit: NASA Scientific Visualization Studio


2015




# Thermal imaging









## Food choices without bees







Sourced online







But they are gardens of trees, not of flowers; green wildernesses. Imagine... for days, even weeks, you must ride with no shade, and the sun overhead, and nothing but the bleached bones of dead animals strewing the track. Then when you come to trees and running water, you will call it a garden. It will not be flowers and their garishness that your eyes crave for, but a green cavern full of shadows and pools where goldfish dart, and the sound of a little stream.

# Useful websites

Green Gecko Studios

- naturalresources.sa.gov.au
- butterflygardening.net.au
- www.npsp.sa.gov.au

## **Useful books**



## The Native Plants of Adelaide

Returning the vanishing natural heritage of the Adelaide Plains to your garden



#### Myoporum species Boobiallas Myoporaceae

Description Ranging from prostrate groundcovers to graceful drooping trees up to 10 m tall, at least five Myoporum species are found on the

Leaves M. plasycarpum (Sugarwood), the only tree in the group, has thin leaves with small serrations near the tip. All the other Myoporums have thick semi-succulent leaves,

Flowers Asymmetrical five-petalled flowers about 1 cm in diameter appear in spring, sometimes downy, often with a pattern of purplish spots on the petals and a delicate scent.

Fruit M. platycarpum and M. parvifolium (the prostrate Cteeping Boobialla) have brown papery capsules. The other shrubby species all develop fleshy berries, ripening either to white or a reddish colour,

Distribution M. acuminatum (Native Myrtle) and M. plasycarpum only occur on the northern Adelaide Plains. M. insulare (Coastal Boobialla) and M. parvifolium are coastal specialists. M. viscosum (Sticky Boobialla) was once found scattered in

Ecology The berries are food for birds.

Similar species The related Eremophila deserti (Turkey Bush) is a mediumsized shrub with similar flowers but is rare on the northern Adelaide Plains.

Cultivation Most species are easily grown from tip cuttings, seeds are often difficult to germinate. Some species can be grown from seed by soaking the stone overnight in warm water, while others are grown by fermenting away the flesh and planting the stone.









The Native Plants of Adelaide



Attracting Butterflies to your Garden

What to Grove and Conserve in the Adelaide Region

Barnerily Gardening Rect

Green Gecko Studios

Chequered Swallowtail

Abundance in Adelaide area: Uncommon Flight: Sep-May Wingspan: in 72 mm; 175 mm Mature larva length: 45 mm

This lovely Swallowtal is an occasional visitor for Activate in the course of as migratury flights, it will breed here when it flight, its catespillar flood plant, but probably carried survive the writers. Young catespillar resemble and dropping during the first flour instars, but in the first are socisty given. Its catespillar food plants (Cultim spp.) make an attractive addition to gardens and butterfiles can also use the flowers as a hoctar source. The Common Grass share burnerly also uses those plants for breading.

interestingly, the butterfly emerges from the pupa in as little as druce weeks or as much as ten months, or longer. This can help the survival of this and adapted species in adverse years.

Caterpillar food plants: Scurl-pess (Culor spp.) Caterpillars eat the flowers and soft green parts of these annual or short-lived perennial plants.

Adetaide native species: tall Scart-pea (Culion australizations: proviously known as Pagraday).

Other South Australian species: Annual Sourt-pos (Cullen chemount), Native Lucerno (Cullen gravaciera), Spreading Sourt-pos (Cullen pallors)

Though comparatively rare in the Adelaloe region, this buttarity still breeds in the area, wherever it encounters the new rare normant examples of its caterpliar bood plant, the Sourt-pea. The Chaquered Swallowtall is one of the largest butterflare regularly seen in southern Australia and as such, with its pair yellow and black colouration and rapid flight, attracts artisation from these who have an interest in butterflare.

This butterily is quite different in appearance from other South Australian species. Restly it is described in the process of the process of

The mature caterpliar is roughly cylindrical and sightly humped towards the head. The body colour ranges from green to yellow groon, with this black areas between the body segments, these are numerous, black-edged, orange spots, and a yellow line along the lower margin of the body.







HOSE ARE SETTLY BUTTER BY JOHN GO HOSE STATE OF TORREST





## Green Gecko Studios

#### **Videos:**

- Marla Spivak TED Talk on 'Why bees are disappearing'.
- Gardening Australia: It comes down to love

#### **Book:**

South Australian Native Bees by Susanne Richards





#### **Movies:**


- Queen of the Sun
- Vanishing of the Bees
- More than Honey
- Messages from bees Japanese documentary (some translation issues)

# Find out more



### **Movies:**

- Food Inc
- Project Wild Thing



Thanks for your bee-autiful work!